Smoke-Free Coalition 3 May 2010

Mississippi Tobacco Data







Cigarette Smoke is a Toxic Air Contaminant

Secondhand tobacco smoke is in the same category as the most toxic automotive and industrial air pollutants

- California EPA

There is No Safe Level of Exposure to Secondhand Smoke

- US Surgeon General

What is secondhand smoke?

Contains more than 4,000 chemical compounds, of which at least 43 are known carcinogens, including:

- Carbon monoxide (auto exhaust)
- Tar (roofing)
- Nicotine (insecticide)
- Acetone (nail polish remover)
- Toluene (industrial solvent)
- Formaldehyde (preservative for dead bodies)
- Phenol (disinfectant)
- Benzene (industrial solvent)
- Benzopyrene (diesel exhaust)
- Hydrogen cyanide (gas chamber poison)
- Arsenic (rat poison)
- DDT (insecticide)

Long Term Harms

- Coronary heart disease
- Lung cancer
- Nasal sinus cancer


Immediate Harms of SHS

- SIDS
- Low birth weight rate
- Ear infections
- Lower respiratory illness in children
- Acute respiratory infections
- More severe asthma
- Impacts cardiovascular system


Smoke-free laws have improved health of workers and patrons

- Improved respiratory health
- Fewer reports of eye, nose, and throat irritation
- Fewer hospital admissions for heart attacks in Pueblo,
 CO and Helena, MT
- Institute of Medicine Report
- Data from Starkville and Hattiesburg hospitals

Heart Attack Admission Decreases in Starkville



Heart Attack Admission Decreases in Hattiesburg



Health and Economic Burden of Secondhand Smoke in the U.S.

Annual U.S. deaths attributable to exposure to secondhand smoke

■ Heart disease: 35,000

■ Lung cancer: 3,000

■ SIDS: 1,900

Estimated annual diseases attributable to exposure to secondhand smoke in the U.S.

- Low birth weight births: 9,700
- Asthma exacerbation in children: 400,000
- Acute lower respiratory illness: 150,000
- Ear infections: 700,000


Estimated direct costs of exposure to secondhand smoke in the U.S., by morbidity

- Lung cancer: \$191 million
- Cervical cancer: \$14 million
- Asthma: \$773 million
- Ear infections: \$53 million
- Chronic pulmonary disease: \$1.2 billion
- Coronary heart disease: \$2.5 billion
- Low birth weight: \$284 million

Public Support 2009 Mississippi Social Climate Survey of Tobacco Control

- 81.7% of Mississippi adults believe that restaurants should be completely smoke free
- 83.6% of Republicans
- 83.1% of Democrats
- 76.1% of Mississippi adults believe that worksites should be completely smoke free
- 80% of Republicans
- 74.5% of Democrat
- Non-partisan support, both Democrats and Republicans support smoke free laws

Mississippi Smoke-Free Communities



Mississippi Population Protected by a Comprehensive Smoke-Free Law

2002	Local Opomances Metcalfe	DATE IMPLEMENTED SEPTEMBER 3RD	% or MS Por. Page 0.04%
2005			
	Oxford	November	
	ABERDEEN	FEBRUARY	
		MARCH 8TH	
	RIDGELAND	July 19th	5.18%
	CORINTH	November	
		November 2nd	
2008			
	Collins	June 8th	
	Cunton		
	LAUREL	December 4th	
	GREENWOOD		
	GRENADA		
	HOLLANDALE		
		Макси 4тн	
	BASSFIELD	MARCH 10th	11.18%
	PRENTISS		
	CRYSTAL SPRINGS		
C	ONTACT OF STREET	Robert McMillen 662.325.27 robert.mcmillen@ssrc. mstobaccodate	12 msstate.edu

Contact

For more information please visit us on the web:

mstobaccodata.org