



UNIVERSITY
REPORT

UNIVERSITY OF
MISSISSIPPI

2019 • mississippi tobacco data



Data Management Coordinator | Emily McClelland, M.S.

Art Director + Graphic Designer | Miranda Robertson

Photographer (Cover Photo) | Robert Jordan



UNIVERSITY SURVEY OF TOBACCO CONTROL

/// UNIVERSITY OF MISSISSIPPI | 2019

ACKNOWLEDGMENT

This survey was made possible through support provided by the Mississippi State Department of Health. The opinions expressed herein are those of the authors and do not necessarily reflect the views of the Mississippi State Department of Health.

The survey instrument design was based on an extensive review of extant instruments such as the Social Climate Survey of Tobacco Control, the Current Population Survey – Tobacco Use Supplement, the National College Health Risk Behavior Survey, the Youth Tobacco Survey, the Harvard School of Public Health College Alcohol Survey, and was supplemented by additional relevant items.

METHODS


Researchers at the Social Science Research Center at Mississippi State University (MSU) administered a web-based survey to undergraduate students at the University of Mississippi (UM). This web-based approach has extraordinary potential as the university provides internet access to students in multiple settings and issues each student an e-mail account. Moreover, students rely on e-mail and the internet to communicate with peers, professors, and university administrators.

Information Technology Services at UM provided 16,220 undergraduate student e-mail addresses. Of these 16,220 students, 8,110 were sampled and 352 participated in the survey. All respondents were 18 years of age

or older and undergraduate students at UM during the Spring 2019 term. Respondents were screened before beginning the survey, and students who were not yet 18 years of age during the Spring 2019 term were not allowed to continue the survey.


Each student in the sample received an e-mail asking for their participation in the survey. Students who chose to participate clicked on an embedded URL address that linked the student to the survey web site. Respondents who did not complete the survey in one session were able to restart the survey at the last question answered. Up to two reminder e-mails were sent requesting the student's continued participation. Two reminder emails were also sent to each student who did not respond to the initial request for participation. No more than three emails were sent to any student in the sample.

Several precautions were taken to protect the privacy of respondents. Personal identifiers (e-mail addresses) were kept separate from the information collected from each respondent. This was accomplished by anonymizing responses so that no personal information was recorded and the contact association was removed.



INDIVIDUAL PRACTICES & EXPOSURE TO TOBACCO SMOKE


2019



- The percentage of University of Mississippi students who reported smoking is not allowed anywhere in their home was significantly higher for females (92.7%) than for males (81.5%), $p < .01$. Juniors (94.1%) were 5.0 times as likely and seniors (91.0%) were 3.2 times as likely as sophomores (76.1%) to report smoking is not allowed anywhere in their home, $p = .01$.
- The percentage of University of Mississippi students who reported they were not exposed to secondhand smoke in a vehicle in the past seven days was significantly higher for females (84.8%) than for males (74.6%), $p = .03$.

SMOKE-FREE POLICIES IN THE COMMUNITY


2019



- The percentage of University of Mississippi students who reported witnessing someone smoking in an indoor public place where smoking is prohibited by law was significantly higher for females (23.7%) than for males (12.3%), $p = .01$.

HEALTH BELIEFS & COUNSELING + MASS MEDIA


2019



- The percentage of University of Mississippi students who reported their doctor or someone at their doctor's office has discussed the dangers of smoking with them was significantly higher for males (27.1%) than for females (24.7%), $p = .04$.


CIGARETTE SMOKING

2019



- The percentage of University of Mississippi students who reported ever trying cigarettes was significantly higher for males (79.0%) than for females (47.8%), $p < .001$.
- The percentage of University of Mississippi students who reported smoking cigarettes in the past thirty days was significantly higher for males (35.0%) than for females (14.5%), $p < .001$.

TOBACCO PRODUCT USE /// ◀ 2019 ▶




- The percentage of University of Mississippi students who reported having ever tried smoking tobacco in a hookah or water pipe was significantly higher for males (41.3%) than for females (26.1%), $p < .01$. Juniors (42.3%) were 2.3 times as likely and seniors (38.2%) were 1.9 times as likely as freshmen (24.2%) to report having ever tried smoking tobacco in a hookah or water pipe, $p < .05$.
- The percentage of University of Mississippi students who reported having ever used an electronic cigarette was significantly higher for males (80.3%) than for females (66.1%), $p < .01$. Freshmen (74.4%) were 2.1 times as likely and sophomores (86.8%) were 4.7 times as likely as seniors (58.4%) to report having ever used an electronic cigarette, $p = .02$.
- The percentage of University of Mississippi students who reported having used an electronic cigarette in the past thirty days was significantly higher for males (51.5%) than for females (33.3%), $p = .001$. Freshmen (47.8%) were 2.1 times as likely and sophomores (51.3%) were 2.4 times as likely as seniors (30.3%) to report having used an electronic cigarette in the past thirty days, $p = .02$.

TOBACCO PRODUCT USE ——— // ——— ◀ 2019 ▶

- The percentage of University of Mississippi students who reported having used an electronic cigarette in the past thirty days and primarily used a pod-based device was significantly higher for males (41.2%) than for females (26.4%), $p < .01$. Freshmen (36.7%) were 2.1 times as likely and sophomores (42.7%) were 2.7 times as likely as seniors (21.3%) to report having used an electronic cigarette and primarily used a pod-based device in the past thirty days, $p = .03$.
- The percentage of University of Mississippi students who reported having used smokeless tobacco in the past thirty days was significantly higher for males (32.9%) than for females (0.5%), $p < .001$.
- The percentage of University of Mississippi students who reported having smoked a little filtered cigar in the past thirty days was significantly higher for males (6.4%) than for females (0.5%), $p < .01$.
- The percentage of University of Mississippi students who reported having smoked medium cigars or cigarillos in the past thirty days was significantly higher for males (15.0%) than for females (3.8%), $p < .001$.
- The percentage of University of Mississippi students who reported having smoked large or premium cigars in the past thirty days was significantly higher for males (15.7%) than for females (0.0%), $p < .001$.
- The percentage of University of Mississippi students who reported having smoked flavored little cigars in the past thirty days was significantly higher for males (5.8%) than for females (0.5%), $p < .01$.

SMOKING CESSATION ——— // ——— ◀ 2019 ▶

Current Smokers Who Have Ever Participated in a Quit Program
 Current Smokers Who Stopped for One Day or More in an Attempt to Quit



CONTACT INFORMATION



Robert McMillen, Ph.D.
Social Science Research Center
Mississippi State University

ROBERT.MCMILLEN@SSRC.MSSTATE.EDU
WWW.SSRC.MSSTATE.EDU
WWW.MSSTATE.EDU

One Research Blvd., Suite 103
Starkville, MS 39759

P: 662.325.7127
F: 662.325.7966

For more information visit

MSTOBACCODATA.ORG



MISSISSIPPI STATE UNIVERSITY DOES NOT DISCRIMINATE ON THE BASIS OF RACE, COLOR, RELIGION, NATIONAL ORIGIN, SEX, AGE, DISABILITY, OR VETERAN STATUS. THIS REPORT IS FUNDED BY A GRANT OF THE MISSISSIPPI STATE DEPARTMENT OF HEALTH.



MISSISSIPPI STATE UNIVERSITY™
SOCIAL SCIENCE RESEARCH CENTER